

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Kasl
Catalog Number: R0544S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10090070
Expiration Date: 09/2021
Storage Temperature: -20°C

Storage Conditions: 500 mM KCl, 20 mM Tris-HCl (pH 7.0), 0.1 mM EDTA, 1mM MgCl2, 50%

Glycerol, 0.10% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0544S/L v2.0

| Kasl Component List    |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0544SVIAL             | Kasl                  | 10081964   | Pass                 |  |
| B7204SVIAL             | CutSmart® Buffer      | 10085423   | Pass                 |  |

| Assay Name/Specification  | Lot # 10090070 |
|---|----------------|
| Blue-White Screening (Terminal Integrity) A sample of LITMUS38i vector linearized with a 10-fold excess of KasI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.   | Pass           |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 5 units of Kasl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.                | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pBR322 DNA with KasI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with KasI.                                    | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pBR322 DNA and a minimum of 5 Units of Kasl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |



R0544S / Lot: 10090070

Page 1 of 2



| Assay Name/Specification   | Lot # 10090070 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE) Kasl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

03 Nov 2020

Michael Tonello

Packaging Quality Control Inspector

03 Nov 2020



R0544S / Lot: 10090070

Page 2 of 2