

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Pacl
Catalog Number: R0547S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pNEB193 DNA in 1 hour at 37°C in a total reaction volume of 50

μl.

Lot Number: 10032193
Expiration Date: 09/2020
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0547S/L v1.0

Pacl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0547SVIAL	Pacl	10020944	Pass	
B7204SVIAL	CutSmart® Buffer	10021125	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10021136	Pass	

Assay Name/Specification	Lot # 10032193
Blue-White Screening (Terminal Integrity)	Pass
A sample of pNEB193 vector linearized with a 10-fold excess of PacI, religated and	
transformed into an E. coli strain expressing the LacZ beta fragment gene results in	
<1% white colonies.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 30 Units of Pacl incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 100 units of Pacl incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pNEB193 DNA with Pacl, ~75% of the DNA fragments	



R0547S / Lot: 10032193

Page 1 of 2

Assay Name/Specification	Lot # 10032193
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Pacl.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pNEB193 DNA and a minimum of 100 units of PacI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Pacl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis
Production Scientist

26 Sep 2018

Michael Tonello

Packaging Quality Control Inspector

03 Jan 2019



R0547S / Lot: 10032193