

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Pacl
Catalog Number: R0547S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pNEB193 DNA in 1 hour at 37°C in a total reaction volume of 50

μl.

Packaging Lot Number: 10093228
Expiration Date: 09/2022
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0547S/L v1.0

| Pacl Component List    |                              |            |                      |  |
|------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description        | Lot Number | Individual QC Result |  |
| R0547SVIAL             | Pacl                         | 10082424   | Pass                 |  |
| B7204SVIAL             | CutSmart® Buffer             | 10091458   | Pass                 |  |
| B7024AVIAL             | Gel Loading Dye, Purple (6X) | 10089393   | Pass                 |  |

| Assay Name/Specification   | Lot # 10093228 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE)  | Pass           |
| Pacl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. |                |
| Non-Specific DNase Activity (16 Hour)  | Pass           |
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pNEB193 DNA and a minimum of |                |
| 100 units of PacI incubated for 16 hours at 37°C results in a DNA pattern free of    |                |
| detectable nuclease degradation as determined by agarose gel electrophoresis.        |                |
| Exonuclease Activity (Radioactivity Release)   | Pass           |
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and      |                |
| double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Pacl incubated for 4 |                |
| hours at 37°C releases <0.1% of the total radioactivity.                             |                |
| Endonuclease Activity (Nicking)  | Pass           |
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and  |                |
| a minimum of 30 Units of PacI incubated for 4 hours at 37°C results in <10%          |                |
| conversion to the nicked form as determined by agarose gel electrophoresis.          |                |



R0547S / Lot: 10093228

Page 1 of 2

| Assay Name/Specification   | Lot # 10093228 |
|--|----------------|
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pNEB193 DNA with Pacl, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Pacl. | Pass           |
| Blue-White Screening (Terminal Integrity) A sample of pNEB193 vector linearized with a 10-fold excess of PacI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.          | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

05 Jan 2021

Michael Tonello

Packaging Quality Control Inspector

05 Jan 2021



R0547S / Lot: 10093228

Page 2 of 2