

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | Pacl |
|------------------------|---|
| Catalog Number: | R0547S |
| Concentration: | 10,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μ g of pNEB193 DNA in 1 hour at 37°C in a total reaction volume of 50 μ l. |
| Packaging Lot Number: | 10115411 |
| Expiration Date: | 02/2023 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA |
| Specification Version: | PS-R0547S/L v1.0 |

| Pacl Component List | | | | |
|---------------------|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0547SVIAL | Pacl | 10097999 | Pass | |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10108731 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10111604 | Pass | |

| Assay Name/Specification | Lot # 10115411 |
|---|----------------|
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of pNEB193 DNA with PacI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PacI. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pNEB193 DNA and a minimum of 100 units of Pacl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of PacI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10115411 |
|--|----------------|
| a minimum of 30 Units of PacI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | |
| Blue-White Screening (Terminal Integrity) A sample of pNEB193 vector linearized with a 10-fold excess of PacI, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies. | Pass |
| Protein Purity Assay (SDS-PAGE) | Pass |
| Pacl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 30 Jul 2021

Josh Hersey

Packaging Quality Control Inspector 30 Jul 2021

