

DISCOVERY 240 County Road
GENUINE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Pmel
Catalog Number: R0560S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10068584
Expiration Date: 12/2021
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0560S/L v1.0

Pmel Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0560SVIAL	Pmel	10060897	Pass	
B7204SVIAL	CutSmart® Buffer	10071077	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10065747	Pass	

Assay Name/Specification	Lot # 10068584
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of Lambda DNA and a minimum of	Pass
10 Units of Pmel incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Blue-White Screening (Terminal Integrity) A sample of pNEB193 vector linearized with a 10-fold excess of Pmel, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart <sup>TM</sup> Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 10 Units of Pmel incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Pmel incubated for 4	Pass



R0560S / Lot: 10068584

Page 1 of 2

Assay Name/Specification	Lot # 10068584
hours at 37°C releases <0.1% of the total radioactivity.	
Ligation and Recutting (Terminal Integrity)  After a 10-fold over-digestion of Lambda DNA with Pmel, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Pmel.	Pass

This product has been tested and shown to be in compliance with all specifications.

Penghua Zhang Production Scientist

13 Apr 2020

Jay Minichiello

Packaging Quality Control Inspector

13 Apr 2020



R0560S / Lot: 10068584

Page 2 of 2