

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Apol   |
|------------------------|--|
| Catalog Number:        | R0566S   |
| Concentration:         | 10,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA in 1 hour at 50°C in a total reaction volume of 50 $\mu$ l. |
| Lot Number:            | 10055569   |
| Expiration Date:       | 07/2021  |
| Storage Temperature:   | -20°C  |
| Storage Conditions:    | 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA  |
| Specification Version: | PS-R0566S/L v1.0   |

| Apol Component List |                       |            |                      |  |
|---------------------|-----------------------|------------|----------------------|--|
| NEB Part Number     | Component Description | Lot Number | Individual QC Result |  |
| R0566SVIAL          | Apol                  | 10045348   | Pass                 |  |
| B7203SVIAL          | NEBuffer™ 3.1         | 10052508   | Pass                 |  |

| Assay Name/Specification  | Lot # 10055569 |
|---|----------------|
| <b>Blue-White Screening (Terminal Integrity)</b><br>A sample of pUC19 vector linearized with a 10-fold excess of Apol, religated and<br>transformed into an E. coli strain expressing the LacZ beta fragment gene results in<br><1% white colonies.   | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 μl reaction in NEBuffer 3.1 containing 1 μg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Apol incubated for 4<br>hours at 50°C releases <0.1% of the total radioactivity.    | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 20-fold over-digestion of lambda DNA with Apol, >95% of the DNA fragments<br>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,<br>>95% can be recut with Apol.   | Pass           |
| <b>Non-Specific DNase Activity (16 Hour)</b><br>A 50 µl reaction in NEBuffer 3.1 containing 1 µg of lambda DNA and a minimum of 100<br>Units of Apol incubated for 16 hours at 50°C results in a DNA pattern free of<br>detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification  | Lot # 10055569 |
|---|----------------|
| Protein Purity Assay (SDS-PAGE)<br>Apol is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue<br>detection. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

Production Scientist 12 Aug 2019

al

Jay Minichiello Packaging Quality Control Inspector 11 Sep 2019

