

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Sapl
Catalog Number: R0569S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10110081
Expiration Date: 03/2023
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 500 µg/ml BSA

Specification Version: PS-R0569S/L v1.0

Sapl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0569SVIAL	Sapl	10101567	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10105817	Pass	
B6004SVIAL	rCutSmart™ Buffer	10108412	Pass	

Assay Name/Specification	Lot # 10110081
Protein Purity Assay (SDS-PAGE)	Pass
SapI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with SapI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with Sapl.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	
30 Units of Sapl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 μl reaction in CutSmart [™] Buffer containing 1 μg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 30 units of SapI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	



R0569S / Lot: 10110081

Page 1 of 2

Assay Name/Specification	Lot # 10110081
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled LITMUS38i DNA	
and a minimum of 10 units of Sapl incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

16 Jun 2021

Michael Tonello

Packaging Quality Control Inspector

16 Jun 2021



R0569S / Lot: 10110081

Page 2 of 2