

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | BseRI   |
|------------------------|---|
| Catalog Number:        | R0581L  |
| Concentration:         | 5,000 U/ml  |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 μg<br>of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl. |
| Packaging Lot Number:  | 10119824  |
| Expiration Date:       | 11/2023   |
| Storage Temperature:   | -20°C   |
| Storage Conditions:    | 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA  |
| Specification Version: | PS-R0581S/L v1.0  |

| BseRI Component List |                       |            |                      |  |
|----------------------|-----------------------|------------|----------------------|--|
| NEB Part Number      | Component Description | Lot Number | Individual QC Result |  |
| R0581LVIAL           | BseRI                 | 10119828   | Pass                 |  |
| B6004SVIAL           | rCutSmart™ Buffer     | 10126633   | Pass                 |  |

| Assay Name/Specification  | Lot # 10119824 |
|---|----------------|
| Exonuclease Activity (Radioactivity Release)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 25 units of BseRI incubated for 4<br>hours at 37°C releases <0.1% of the total radioactivity. | Pass           |
| Non-Specific DNase Activity (16 Hour)<br>A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of<br>5 Units of BseRI incubated for 16 hours at 37°C results in a DNA pattern free of<br>detectable nuclease degradation as determined by agarose gel electrophoresis.           | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 10-fold over-digestion of Lambda DNA with BseRI, >95% of the DNA fragments<br>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,<br>~75% can be recut with BseRI.   | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Penghua Zhang

Penghua Zhang Production Scientist 16 Nov 2021

Michae 11.

Michael Tonello Packaging Quality Control Inspector 16 Nov 2021

