

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | NspI  |
|------------------------|---|
| Catalog #:             | R0602S/L  |
| Concentration:         | 10,000 units/ml   |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in 1 hour at 37°C in a total reaction<br>volume of 50 μl. |
| <i>Lot</i> #:          | 0101405   |
| Assay Date:            | 05/2014   |
| Expiration Date:       | 5/2016  |
| Storage Temp:          | -20 °C  |
| Storage Conditions:    | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 μg/ml<br>BSA                               |
| Specification Version: | PS-R0602S/L v1.0  |
| Effective Date:        | 24 Apr 2014   |

| Assay Name/Specification (minimum release criteria)  | Lot #0101405 |
|--|--------------|
| <b>Endonuclease Activity (Nicking)</b> - A 50 μl reaction in CutSmart <sup>™</sup> Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 10 units of NspI incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.                             | Pass         |
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 $\mu$ l reaction in CutSmart <sup>TM</sup> Buffer containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 50 units of NspI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass         |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with NspI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NspI.   | Pass         |
| <b>Non-Specific DNase Activity (16 Hour)</b> - A 50 $\mu$ l reaction in CutSmart <sup>TM</sup> Buffer containing 1 $\mu$ g of Lambda DNA and a minimum of 50 Units of NspI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.     | Pass         |
| <b>Protein Purity Assay (SDS-PAGE)</b> - NspI is $\geq$ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.  | Pass         |

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

otunen

Authorized by Derek Robinson 24 Apr 2014



. Her

Inspected by David Hough 24 Apr 2014