

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Sfol
Catalog Number: R0606S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 μg

of Lambda DNA (HindIII digest) in rCutSmart Buffer in 1 hour at 37°C

in a total reaction volume of 50 μl.

Packaging Lot Number: 10167911
Expiration Date: 10/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

200 μg/ml rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R0606S/L v2.0

Sfol Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0606SVIAL	Sfol	10167910	Pass	
B6004SVIAL	rCutSmart™ Buffer	10164465	Pass	

Assay Name/Specification	Lot # 10167911
Protein Purity Assay (SDS-PAGE)	Pass
Sfol is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled LITMUS28i DNA and a minimum of 30 units of Sfol incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of SfoI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 30 units of SfoI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass



R0606S / Lot: 10167911

Page 1 of 2

Assay Name/Specification	Lot # 10167911
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda-HindIII DNA with SfoI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SfoI.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun

Production Scientist

25 Oct 2022

Michael Tonello

Packaging Quality Control Inspector

31 Oct 2022



R0606S / Lot: 10167911

Page 2 of 2