

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	BtsI-v2
Catalog Number:	R0667L
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μ g of Lambda DNA in 1 hour at 55°C in a total reaction volume of 50 μ l.
Packaging Lot Number:	10091743
Expiration Date:	08/2022
Storage Temperature:	-20°C
Storage Conditions:	50 mM KCl , 10 mM Tris-HCl, 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 200 μg/ml BSA (pH 7.4 @ 25°C)
Specification Version:	PS-R0667S/L v2.0

BtsI-v2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0667LVIAL	BtsI-v2	10080858	Pass	
B7204SVIAL	CutSmart® Buffer	10085425	Pass	

Assay Name/Specification	Lot # 10091743
Protein Purity Assay (SDS-PAGE) BtsI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 10 units of BtsI-v2 incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BtsI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with BtsI-v2.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of BtsI-v2 incubated for 15 minutes at 55°C results in complete digestion as determined by agarose gel electrophoresis.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10091743
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 50 units of BtsI-v2 incubated for	
4 hours at 55°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Pendina Zhang

Penghua Zhang Production Scientist 20 Nov 2020

Michae

Michael Tonello Packaging Quality Control Inspector 20 Nov 2020

