

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	BsrFI-v2
Catalog Number:	R0682S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of pBR322 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.
Lot Number:	10053565
Expiration Date:	08/2020
Storage Temperature:	-20°C
Storage Conditions:	250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton® X-100, 200 μg/ml BSA, (pH 7.4 @ 25°C)
Specification Version:	PS-R0682S/L v2.0

BsrFI-v2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0682SVIAL	BsrFI-v2	10053564	Pass	
B7204SVIAL	CutSmart® Buffer	10046087	Pass	

Assay Name/Specification	Lot # 10053565
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 30 units of BsrFI-v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of pBR322 DNA and 1 µl of BsrFI-v2 incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) BsrFI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of pBR322 DNA and a minimum of 10 units of BsrFI-v2 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass





be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10053565
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of pBR322 DNA with BsrFI-v2, >95% of the DNA	
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
fragments, >95% can be recut with BsrFI-v2.	

This product has been tested and shown to be in compliance with all specifications.

Production Scientist 09 Aug 2019

204-

Minichiello Packaging Quality Control Inspector 21 Aug 2019

