

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: BspQI
Catalog Number: R0712S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 50°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10201883
Expiration Date: 04/2025
Storage Temperature: -20°C

Storage Conditions: 500 mM KCl , 20 mM Tris-HCl (pH 7.0), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, 0.10 % TritonX-100, 500 µg/ml BSA

Specification Version: PS-R0712S/L v2.0

BspQI Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0712SVIAL	BspQl	10185368	Pass	
B6003SVIAL	NEBuffer™ r3.1	10182164	Pass	

Assay Name/Specification	Lot # 10201883
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled M13mp18 DNA and a minimum of 10 units of BspQl incubated for 4 hours at 50°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of BspQl incubated for 4 hours at 50°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BspQI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BspQI.	Pass
Non-Specific DNase Activity (16 hour) A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 10 Units of BspQl incubated for 16 hours at 50°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE:	Pass



R0712S / Lot: 10201883

Page 1 of 2

Assay Name/Specification	Lot # 10201883
although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.	
Protein Purity Assay (SDS-PAGE) BspQI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun Production Scientist 05 Apr 2023

Michael Tonello

Packaging Quality Control Inspector

10 Aug 2023



R0712S / Lot: 10201883