

## New England Biolabs Certificate of Analysis

**Product Name:** PaqCI<sup>®</sup>  
**Catalog Number:** R0745L  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA incubated for 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10156415  
**Expiration Date:** 01/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0745S/L v1.0

PaqCI <sup>®</sup> Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
S0532LVIAL	PaqCI <sup>®</sup> Activator	10156991	Pass
R0745LVIAL	PaqCI <sup>®</sup>	10156414	Pass
B6004SVIAL	rCutSmart <sup>™</sup> Buffer	10153337	Pass

Assay Name/Specification	Lot # 10156415
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of Lambda DNA and 1 µl of PaqCI <sup>®</sup> incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> PaqCI <sup>®</sup> is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 ul reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of PaqCI <sup>®</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of Lambda DNA and a minimum of 30 units of PaqCI <sup>®</sup> incubated for 16 hours at 37°C results in a DNA pattern free of	Pass

Assay Name/Specification	Lot # 10156415
detectable nuclease degradation as determined by agarose gel electrophoresis.	
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with PaqCI<sup>®</sup> , &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with PaqCI<sup>®</sup> .</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart<sup>®</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 units of PaqCI<sup>®</sup> incubated for 4 hours at 37°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
14 Jul 2022



Josh Hersey  
Packaging Quality Control Inspector  
14 Jul 2022