

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Bcll-HF®
Catalog Number:	R3160L
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of 50 μl.
Lot Number:	10019564
Expiration Date:	08/2020
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 500 μg/ml BSA, (pH 7.4 @ 25°C)
Specification Version:	PS-R3160S/L v1.0

Bcll-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3160LVIAL	Bcll-HF®	10019163	Pass	
B7204SVIAL	CutSmart® Buffer	10015394	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10013725	Pass	

Assay Name/Specification	Lot # 10019564
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of BclI-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda dam- DNA and 1 µl of Bcll-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda dam- DNA with BcII-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BcII-HF.	Pass
Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda dam- DNA and a	Pass





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10019564
minimum of 60 units of BcII-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) Bcll-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

You

Tony Spear-Alfonso Production Scientist 08 Aug 2018

Josh Hersey Packaging Quality Control Inspector 17 Aug 2018

