

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Bcll-HF®
Catalog Number:	R3160S
Concentration:	20,000 U/mI
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 $\mu$ g of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of 50 $\mu$ l.
Packaging Lot Number:	10091879
Expiration Date:	12/2022
Storage Temperature:	-20°C
Storage Conditions:	300 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 500 μg/ml BSA, (pH 7.4 @ 25°C)
Specification Version:	PS-R3160S/L v1.0

Bcll-HF® Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R3160SVIAL	BclI-HF®	10091881	Pass	
B7204SVIAL	CutSmart® Buffer	10091032	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10089401	Pass	

Assay Name/Specification	Lot # 10091879
Protein Purity Assay (SDS-PAGE) Bcll-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda dam- DNA and a minimum of 60 units of BcII-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda dam- DNA with BclI-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BclI-HF.	Pass
Functional Testing (15 minute Digest) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda dam- DNA and 1 μl of	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10091879
Bcll-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and	
double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Bcll-HF incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Penghua Zhang Production Scientist 16 Dec 2020

Michae 11.6

Michael Tonello Packaging Quality Control Inspector 16 Dec 2020

