

be INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

| Product Name: | BstEll-HF® |
|------------------------|--|
| Catalog Number: | R3162M |
| Concentration: | 100,000 U/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 μ g of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μ l. |
| Lot Number: | 10008872 |
| Expiration Date: | 05/2020 |
| Storage Temperature: | -20°C |
| Storage Conditions: | 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA |
| Specification Version: | PS-R3162M v1.0 |

| BstEll-HF® Component List | | | | |
|---------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R3162MVIAL | BstEII-HF® | 10008886 | Pass | |
| B7204SVIAL | CutSmart® Buffer | 10013537 | Pass | |
| B7024SVIAL | Gel Loading Dye, Purple (6X) | 0261805 | Pass | |

| Assay Name/Specification | Lot # 10008872 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 units of BstEII-HF™ incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 60 units of BstEII-HF [™] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BstEII-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BstEII-HF™. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 60 units of BstEII-HF™ incubated for 16 hours at 37°C results in a DNA pattern free | Pass |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification | Lot # 10008872 |
|--|----------------|
| of detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist 23 May 2018

Michae m. //

Michael Tonello Packaging Quality Control Inspector 31 Jul 2018

