

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	SbfI-HF®
Catalog Number:	R3642L
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1μg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10145338
Expiration Date:	03/2024
Storage Temperature:	-20°C
Storage Conditions:	200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Specification Version:	PS-R3642S/L v1.0

SbfI-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3642LVIAL	SbfI-HF®	10140489	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10138405	Pass	
B6004SVIAL	rCutSmart™ Buffer	10138403	Pass	

Assay Name/Specification	Lot # 10145338
Protein Purity Assay (SDS-PAGE) SbfI-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 20 Units of SbfI-HF <sup>™</sup> incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of SbfI-HF <sup>™</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with SbfI-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SbfI-HF™.	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10145338
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and a minimum of 20 Units of Sbfl-HF <sup>™</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Peperlog

Penghua Zhang Production Scientist 29 Mar 2022

Mid

Michael Tonello Packaging Quality Control Inspector 29 Mar 2022

