

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: SbfI-HF®
Catalog Number: R3642L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 1015925
Expiration Date: 08/2024
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R3642S/L v1.0

SbfI-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3642LVIAL	SbfI-HF®	10159250	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10156431	Pass	
B6004SVIAL	rCutSmart™ Buffer	10156433	Pass	

Assay Name/Specification	Lot # 10159251
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 20 Units of Sbfl-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with SbfI-HF [™] , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SbfI-HF [™] .	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Sbfl-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of supercoiled pBR322 DNA and a minimum of 20 Units of Sbfl-HF™ incubated for 4 hours at 37°C results in <20%	Pass



R3642L / Lot: 10159251

Page 1 of 2

Assay Name/Specification	Lot # 10159251
conversion to the nicked form as determined by agarose gel electrophoresis.	
Protein Purity Assay (SDS-PAGE) Sbfl-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

29 Aug 2022

Erin Varney

Packaging Quality Control Inspector

29 Aug 2022



R3642L / Lot: 10159251

Page 2 of 2