

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bsal-HF®v2
Catalog Number: R3733L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number:10030552Expiration Date:09/2020Storage Temperature:-20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA,

50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-R3733S/L v1.0

| Bsal-HF®v2 Component List | | | | |
|---------------------------|------------------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R3733LVIAL | Bsal-HF®v2 | 10023371 | Pass | |
| B7204SVIAL | CutSmart® Buffer | 10021120 | Pass | |
| B7024SVIAL | Gel Loading Dye, Purple (6X) | 10021129 | Pass | |

| Assay Name/Specification | Lot # 10030552 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 units of Bsal-HF®v2 incubated for 4 hours at 37°C results in <20% | Pass |
| conversion to the nicked form as determined by agarose gel electrophoresis. | |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Bsal-HF®v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with Bsal-HF®v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Bsal-HF®v2. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 μl reaction in CutSmart® Buffer containing 1 μg of pXba DNA and a minimum of 60 units of Bsal-HF®v2 incubated for 16 hours at 37°C results in a DNA pattern free of | Pass |



R3733L / Lot: 10030552

Page 1 of 2



| Assay Name/Specification | Lot # 10030552 |
|---|----------------|
| detectable nuclease degradation as determined by agarose gel electrophoresis. | |
| Protein Purity Assay (SDS-PAGE) Bsal-HF®v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso **Production Scientist**

02 Nov 2018

Michael Tonello

Packaging Quality Control Inspector

28 Nov 2018

