

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Bsal-HF®v2
Catalog Number: R3733L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10068115
Expiration Date: 02/2022
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA,

50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-R3733S/L v1.0

Bsal-HF®v2 Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R3733LVIAL	Bsal-HF®v2	10067046	Pass	
B7204SVIAL	CutSmart® Buffer	10064411	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10059230	Pass	

Assay Name/Specification	Lot # 10068115
Protein Purity Assay (SDS-PAGE) Bsal-HF®v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	Pass
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of pXba DNA and a minimum of 60 units of Bsal-HF®v2 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pXba DNA with Bsal-HF®v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Bsal-HF®v2.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Bsal-HF®v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass



R3733L / Lot: 10068115 Page 1 of 2

Assay Name/Specification	Lot # 10068115
Endonuclease Activity (Nicking) A 50 μl reaction in CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 units of Bsal-HF®v2 incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Anthony Francis
Production Scientist

12 Feb 2020

Jay Minichiello

Packaging Quality Control Inspector

27 Feb 2020

