

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Bsal-HF®v2
Catalog Number: R3733L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10132318
Expiration Date: 07/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA,

50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-R3733S/L v1.0

Bsal-HF®v2 Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R3733LVIAL	Bsal-HF®v2	10116343	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10126634	Pass	
B6004SVIAL	rCutSmart™ Buffer	10127377	Pass	

Assay Name/Specification	Lot # 10132318
Protein Purity Assay (SDS-PAGE) Bsal-HF®v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	Pass
detection.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and	Pass
double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Bsal-HF®v2 incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Bsal-HF®v2 incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of pXba DNA with Bsal-HF®v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
fragments, >95% can be recut with Bsal-HF®v2.	



R3733L / Lot: 10132318

Page 1 of 2

Assay Name/Specification	Lot # 10132318
Non-Specific DNase Activity (16 Hour)	Pass
A 50 μl reaction in CutSmart® Buffer containing 1 μg of pXba DNA and a minimum of 60	
units of Bsal-HF®v2 incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

10 Dec 2021

Michael Tonello

Packaging Quality Control Inspector

10 Dec 2021



R3733L / Lot: 10132318

Page 2 of 2