

DISCOVERY 240 County Road
ENUINE Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bsal-HF®v2
Catalog Number: R3733S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10121327
Expiration Date: 07/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 200 µg/ml BSA,

50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-R3733S/L v1.0

Bsal-HF®v2 Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3733SVIAL	Bsal-HF®v2	10116344	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10121393	Pass	
B6004SVIAL	rCutSmart™ Buffer	10121395	Pass	

Assay Name/Specification	Lot # 10121327
Protein Purity Assay (SDS-PAGE) Bsal-HF®v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue	Pass
detection.	
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with Bsal-HF®v2, >95% of the DNA	Pass
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated	
fragments, >95% can be recut with Bsal-HF®v2.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of pXba DNA and a minimum of 60 units of Bsal-HF®v2 incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Bsal-HF®v2 incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	



R3733S / Lot: 10121327

Page 1 of 2

Assay Name/Specification	Lot # 10121327
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 100 units of Bsal-HF®v2 incubated	
for 4 hours at 37°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

05 Oct 2021

Josh Hersey

Packaging Quality Control Inspector

05 Oct 2021

