240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: Salt-T4® DNA Ligase

Catalog #: M0467S/L

Concentration: 400,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to give 50% ligation of 6 µg of Lambda-HindIII DNA in 30 minutes

at 25°C in a total reaction volume of 20 µl in 1X T4 DNA Ligase Reaction Buffer supplemented with 100 mM NaCl.

Shelf Life: 24 months
Storage Temp: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0467S/L v2.0

Effective Date: 17 Apr 2023

Assay Name/Specification (minimum release criteria)

DNase Activity (Labeled Oligo, 3' extension) - A 50 μ l reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 2000 units of Salt-T4® DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.

DNase Activity (Labeled Oligo, 5' extension) - A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 2000 units of Salt-T4® DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.

Double Stranded DNase Activity (Labeled Oligo) - A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 2000 units of Salt-T4® DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.

Endonuclease Activity (Nicking) - A 50 μl reaction in NEBuffer 1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 400 units of Salt-T4® DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in NEBuffer 1 containing 1 μ g of CIP-treated Lambda-HindIII DNA and a minimum of 400 units of Salt-T4® DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - Salt-T4® DNA Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)

qPCR DNA Contamination (*E. coli* Genomic) - A minimum of 400 units of Salt-T4® DNA Ligase is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.

RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Salt-T4® DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Single Stranded DNase Activity (FAM-Labeled Oligo) - A 50 μ l reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 2000 units of Salt-T4® DNA Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Namy Consolm

Date 17 Apr 2023

Nancy Considine Quality Approver





