240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: EnGen® Spy Cas9 NLS

Catalog #: M0646T/M

Concentration: $20 \mu M$ Unit Definition: N/AShelf Life: 24 months Storage Temp: -20°C

Storage Conditions: 10 mM Tris-HCl , 300 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @, 25°C)

Specification Version: PS-M0646T/M v2.0

Effective Date: 11 Jul 2018

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 μl reaction in NEBuffer 3.1 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in NEBuffer 3.1 containing 1 μg of a mixture of single and doublestranded [³H] E. coli DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (Targeted Digestion) - A 20 µl reaction in NEBuffer 3.1 containing 20 nM of 100 bp FAM and ROX-labeled double-stranded target DNA, 100 nM sgRNA, and 100 nM EnGen® Spy Cas9 NLS incubated for 1 hour at 37°C results in ≥90% targeted digestion of the substrate DNA as determined by capillary electrophoresis.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - EnGen® Spy Cas9 NLS is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 picomole of EnGen® Spy Cas9 NLS is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Derek Robinson

Director of Quality Control







Date