240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

21 May 2013

Date

www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name: PacI

Catalog #: R0547S/L

Concentration: 10,000 units/ml

Unit Definition:

One unit is defined as the amount of enzyme required to digest 1 µg of pNEB193 DNA in 1 hour at 37°C in a total reaction

volume of 50  $\mu$ l.

Shelf Life: 24 months Storage Temp:  $-20 \, ^{\circ}\text{C}$ 

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA

Specification Version: PS-R0547S/L v1.0

Effective Date: 21 May 2013

## Assay Name/Specification (minimum release criteria)

**Blue-White Screening (Terminal Integrity)** - A sample of pNEB193 vector linearized with a 10-fold excess of PacI, religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in CutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of supercoiled PhiX174 DNA and a minimum of 30 Units of PacI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50  $\mu$ l reaction in CutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of a mixture of single and double-stranded [  $^3$ H] *E. coli* DNA and a minimum of 100 units of PacI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of pNEB193 DNA with PacI,  $\sim$ 75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PacI.

Non-Specific DNase Activity (16 Hour) - A 50 μl reaction in CutSmart<sup>TM</sup> Buffer containing 1 μg of pNEB193 DNA and a minimum of 100 units of PacI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - PacI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.

Derek Robinson

Director of Quality Control







<sup>\*</sup> The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.