240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: AsiSI

Catalog #: R0630S/L/V
Concentration: 10,000 units/ml

Unit Definition:

One unit is defined as the amount of enzyme required to digest 1 µg of XhoI digested pXba in 1 hour at 37°C in a total

reaction volume of 50 µl.

Shelf Life: 18 months
Storage Temp: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA

Specification Version: PS-R0630S/L v1.0

Effective Date: 08/21/2013

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in CutSmartTM Buffer containing 1 μg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 10 units of AsiSI incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 2-fold over-digestion of pXbaI (Xho digested) DNA with AsiSI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AsiSI.

Non-Specific DNase Activity (16 hour) - A 50 µl reaction in CutSmartTM Buffer containing 1 µg of pXbaI (Xho digested) DNA and a minimum of 10 Units of AsiSI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Derek Robinson Quality Approver







08/21/2013

Date