

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name:	SalI-HF®
Catalog #:	R3138S/L
Concentration:	20,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 μl.
Shelf Life:	24 months
Storage Temp:	-20 °C
Storage Conditions:	50 mM KCl, 10 mM Tris-HCl (pH 7.5), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 300 μg/ml BSA
Specification Version:	PS-R3138S/L v1.0
Effective Date:	01 May 2013

Assay Name/Specification (minimum release criteria)

**Blue-White Screening (Terminal Integrity)** - A sample of pUC19 vector linearized with a 10-fold excess of SalI-HF<sup>TM</sup>, religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in CutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of supercoiled PhiX174 DNA and a minimum of 20 Units of SalI-HF<sup>TM</sup> incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50  $\mu$ l reaction in CutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of a mixture of single and doublestranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 200 units of SalI-HF<sup>TM</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Ligation and Recutting (Terminal Integrity)** - After a 50-fold over-digestion of Adenovirus-2 DNA with SalI-HF<sup>TM</sup>, >95% of the DNA fragments can be ligated with T4 DNA ligase in 4 hours at 25°C. Of these ligated fragments, >95% can be recut with SalI-HF<sup>TM</sup>.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart<sup>™</sup> Buffer containing 1 µg of pBR322 DNA and a minimum of 200 Units of SalI-HF<sup>™</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

Derek Robinson Director of Quality Control



Date 01 May 2013