240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

10 Jun 2013

Date

www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: NcoI-HF®
Catalog #: R3193M

Concentration: 100,000 units/ml

Unit Definition:

One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction

volume of 50 μ l.

Shelf Life: 24 months Storage Temp: $-20 \, ^{\circ}$ C

Storage Conditions: 200 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA

Specification Version: PS-R3193M v1.0
Effective Date: 10 Jun 2013

Assay Name/Specification (minimum release criteria)

Blue-White Screening (Terminal Integrity) - A sample of LITMUS28i vector linearized with a 10-fold excess of NcoI-HFTM, religated and transformed into an $E.\ coli$ strain expressing the LacZ beta fragment gene results in <1% white colonies.

Endonuclease Activity (Nicking) - A 50 μl reaction in CutSmartTM Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 Units of NcoI-HFTM incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in CutSmartTM Buffer containing 1 μ g of a mixture of single and double-stranded [3 H] *E. coli* DNA and a minimum of 100 units of NcoI-HFTM incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 20-fold over-digestion of Lambda DNA with NcoI-HFTM, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NcoI-HFTM.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of NcoI-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - NcoI-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

Derek Robinson

Director of Quality Control







^{*} The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.