240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: Control LAMP Primer Mix (rActin)

Catalog #: S0164B

Concentration: 10X Concentrate

Shelf Life: 24 months

Storage Temp: -20°C

Composition (1X): Proprietary

Specification Version: PS-S0164B v2.0

Effective Date: 03 Mar 2022

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 5 μ l of Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Functional Testing (LAMP primers) - A 25 μ l reaction in 1X WarmStart® LAMP Master Mix with UDG in the presence of LAMP Fluorescent Dye and 1X Control LAMP Primer Mix (rActin) containing 10 ng human RNA results in a threshold time of \leq 20 minutes as determined by fluorescent detection. Reactions that lack human RNA template remain negative over a 30 minute incubation at 65°C.

Non-Specific DNase Activity (16 Hour) - A 50 μl reaction in NEBuffer 2 containing 1 μg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 μl of Control LAMP Primer Mix (rActin) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Phosphatase Activity (pNPP) - A 200 μ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 μ l of Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Control LAMP Primer Mix (rActin) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Derek Robinson

Director, Quality Control





