

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name:	Goat Anti-Rat IgG Magnetic Beads
Catalog #:	<i>\$1433\$</i>
Concentration:	20 mg/ml
Shelf Life:	36 months
Storage Temp:	4°C
Storage Conditions:	0.02 % NaN3, 0.05 % Tween® 20, 0.1 % BSA, 1 X PBS, (pH 7.4 @ 25°C)
Specification Version:	PS-S1433S v2.0
Effective Date:	25 Oct 2021

Assay Name/Specification (minimum release criteria)

**Binding Capacity (Magnetic Beads)** - Goat Anti-Rat IgG Magnetic Beads (1 mg) were equilibrated and incubated with 200 µl of >0.5 mg/ml Purified Rat IgG for 1 hour at 25°C, then washed and the Rat IgG eluted. Binding capacity was determined to be >5 µg of Rat IgG per mg of beads.

Functional Binding Assay (Qualitative) - Goat Anti-Rat IgG Magnetic Beads (1 mg) were equilibrated and incubated with 500 µl of Serum containing Rat IgG for 1 hour at 25°C, then washed, eluted and evaluated by Tris-Glycine gel to confirm low non-specific binding of extract proteins and high isolation of target.

Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in Goat Anti-Rat IgG Magnetic Bead Storage Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**RNase Activity (Extended Digestion)** - A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1  $\mu$ l of Goat Anti-Rat IgG Magnetic Beads is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit <u>www.neb.com/trademarks</u> for additional information.

Date 25 Oct 2021

Derek Robinson Director, Quality Control



PS-S1433S v2.0 Page 1 of 1