

New England Biolabs Certificate of Analysis

Product Name: rCutSmart™ Buffer
Catalog Number: B6004S
Concentration: 10 X Concentrate
Packaging Lot Number: 10148896
Expiration Date: 04/2025
Storage Temperature: -20°C
Specification Version: PS-B6004S/V v1.0
Composition (1X): 50 mM Potassium Acetate, 20 mM Tris Acetate, 10 mM Magnesium Acetate, 100 µg/ml rAlbumin, (pH 7.9 @ 25°C)

rCutSmart™ Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B6004SVIAL	rCutSmart™ Buffer	10144739	Pass

Assay Name/Specification	Lot # 10148896
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of MscI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 unit of EcoRI-HF® incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Functional Testing (Restriction Digest, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and 1 unit of ClaI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
Conductivity (buffers/solutions) The conductivity of 10X rCutSmart™ Buffer is between 40 and 46 mS/cm at 25°C.	Pass
RNase Activity (Buffer) A 10 µl reaction in 1X rCutSmart™ Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent	Pass


Assay Name/Specification	Lot # 10148896
detection.	
<p>Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 1X rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>pH (buffers/solutions) The pH of 10X rCutSmart™ Buffer is between pH 7.8 and 8.0 at 25°C.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ben Penta
Production Scientist
25 Apr 2022



Erin Varney
Packaging Quality Control Inspector
25 Apr 2022