

New England Biolabs Certificate of Analysis

Product Name: *SHuffle® T7 Express Competent E.coli*
 Catalog Number: C3029J
 Packaging Lot Number: 10162204
 Expiration Date: 08/2023
 Storage Temperature: -80°C
 Specification Version: PS-C3029J v1.0

SHuffle® T7 Express Competent E.coli Component List

| NEB Part Number | Component Description | Lot Number | Individual QC Result |
|-----------------|--------------------------------------|------------|----------------------|
| C3029JVIAL | SHuffle® T7 Express Competent E.coli | 10151746 | Pass |

| Assay Name/Specification | Lot # 10162204 |
|---|----------------|
| Antibiotic Resistance (Streptomycin) 15 µl of untransformed SHuffle® T7 Express Competent E. coli streaked onto a LB or Rich Broth plate containing Streptomycin will form colonies after incubation for 16 hours at 37°C. | Pass |
| Antibiotic Sensitivity (Chloramphenicol) 15 µl of untransformed SHuffle® T7 Express Competent E. coli streaked onto a LB or Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C. | Pass |
| Antibiotic Sensitivity (Ampicillin) 15 µl of untransformed SHuffle® T7 Express Competent E. coli streaked onto a LB or Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C. | Pass |
| Antibiotic Resistance (Spectinomycin) 15 µl of untransformed SHuffle® T7 Express Competent E. coli streaked onto a LB or Rich Broth plate containing Spectinomycin will form colonies after incubation for 16 hours at 37°C. | Pass |
| Transformation Efficiency 50 µl of SHuffle® T7 Express Competent E. coli cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10 ⁷ cfu/µg of DNA. | Pass |
| Phage Resistance (φ 80) | Pass |


| Assay Name/Specification | Lot # 10162204 |
|--|----------------|
| <p>15 µl of untransformed Shuffle® T7 Express Competent E. coli streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.</p> | |
| <p>Functional Testing (Disulfide Bond Formation) The nuclease NucA requires disulfide bonds for its stability. When expressed at 37°C in E. coli, NucA is toxic to cells only in its oxidized disulfide-bonded state. Transformation of SHuffle® T7 Express Competent E. coli using 100 pg of plasmid that expresses a MBP-NucA fusion results in < 1% of the colonies when compared to a control transformation of its wild type parent strain NEB 10-beta.</p> | Pass |
| <p>Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed Shuffle® T7 Express Competent E. coli streaked onto a LB or Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.</p> | Pass |
| <p>Antibiotic Sensitivity (Kanamycin) 15 µl of untransformed Shuffle® T7 Express Competent E. coli streaked onto a LB or Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p> | Pass |
| <p>Antibiotic Resistance (Nitrofurantoin) 15 µl of untransformed Shuffle® T7 Express Competent E. coli streaked onto a LB or Rich Broth plate containing Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Qiuting Ren
Production Scientist
17 Aug 2022



Anna Sorensen
Packaging Quality Control Inspector
17 Aug 2022