

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | NEB® Stable Competent E. coli (High Efficiency) |
|------------------------|---|
| Catalog Number:        | C3040I  |
| Packaging Lot Number:  | 10167752  |
| Expiration Date:       | 09/2023   |
| Storage Temperature:   | -80°C   |
| Specification Version: | PS-C3040H/I v1.0                                |

| NEB® Stable Competent E. coli (High Efficiency) Component List |   |            |                      |  |
|--|---|------------|----------------------|--|
| NEB Part Number  | Component Description                           | Lot Number | Individual QC Result |  |
| N3041AVIAL   | pUC19 Vector                                    | 10160822   | Pass                 |  |
| C3040IVIAL   | NEB® Stable Competent E. coli (High Efficiency) | 10151517   | Pass                 |  |
| B9035SVIAL   | NEB® 10-beta/Stable Outgrowth Medium            | 10152405   | Pass                 |  |

| Assay Name/Specification   | Lot # 10167752 |
|--|----------------|
| Blue-White Screening ( $\alpha$ -complementation, Competent Cells)<br>NEB® Stable Competent E. coli (High Efficiency) were shown to be suitable for<br>blue/white screening by $\alpha$ -complementation of the $\beta$ -galactosidase gene using pUC19.                                 | Pass           |
| <b>Phage Resistance (<math>\phi</math> 80)</b><br>15 µl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate does not support plaque formation by phage $\phi$ 80 after incubation<br>for 16 hours at 37°C.                              | Pass           |
| <b>Transformation Efficiency</b><br>50 µl of NEB® Stable Competent E. coli (High Efficiency) cells were transformed with<br>100 pg of pUC19 DNA using the transformation protocol provided. Incubation<br>overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e9 cfu/µg of DNA. | Pass           |
| Antibiotic Sensitivity (Ampicillin)<br>15 µl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate containing Ampicillin will not form colonies after incubation for<br>16 hours at 37°C.   | Pass           |
| Antibiotic Sensitivity (Chloramphenicol)<br>15 µl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate containing Chloramphenicol will not form colonies after<br>incubation for 16 hours at 37°C.                                       | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification   | Lot # 10167752 |
|--|----------------|
| <b>Antibiotic Resistance (Tetracycline)</b><br>15 μl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate containingTetracycline will form colonies after incubation for 16<br>hours at 37°C.    | Pass           |
| Antibiotic Resistance (Streptomycin)<br>15 µl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate containing Streptomycin will form colonies after incubation for<br>16 hours at 37°C.          | Pass           |
| Antibiotic Sensitivity (Spectinomycin)<br>15 µl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate containing Spectinomycin will not form colonies after incubation<br>for 16 hours at 37°C.   | Pass           |
| <b>Antibiotic Sensitivity (Kanamycin)</b><br>15 µl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate containing Kanamycin will not form colonies after incubation for<br>16 hours at 37°C.    | Pass           |
| Antibiotic Sensitivity (Nitrofurantoin)<br>15 µl of untransformed NEB® Stable Competent E. coli (High Efficiency) streaked onto<br>a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation<br>for 16 hours at 37°C. | Pass           |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

hh

Lixin An Production Scientist 15 Jul 2022

Anna Sorensen Packaging Quality Control Inspector 17 Oct 2022

