

www.neb.com info@neb.com



## New England Biolabs Certificate of Analysis

Product Name: Taql Methyltransferase

Catalog Number: M0219S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg

Lambda DNA in 1 hour at 65°C in a total reaction volume of 20 μl

against cleavage by Taql restriction endonuclease.

Packaging Lot Number: 10225044
Expiration Date: 01/2026
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

200 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-M0219S v1.0

| Taql Methyltransferase Component List |                              |            |                      |  |
|---------------------------------------|------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                | <b>Component Description</b> | Lot Number | Individual QC Result |  |
| M0219SVIAL                            | Taql Methyltransferase       | 10225043   | Pass                 |  |
| B9003SVIAL                            | S-adenosylmethionine (SAM)   | 10210241   | Pass                 |  |
| B6004SVIAL                            | rCutSmart™ Buffer            | 10224150   | Pass                 |  |

| Assay Name/Specification  | Lot # 10225044 |
|---|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 2 containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of Taql Methyltransferase incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.  | Pass           |
| Functional Testing (Methyltransferase) A 20 μl reaction in CutSmart® Buffer supplemented with 80 μM SAM containing 1 μg of Lambda DNA and 1 unit of Taql Methyltransferase incubated for 1 hour at 65°C followed by heat inactivation results in ≥ 95% protection from digestion with 10 units of Taql in CutSmart® Buffer incubated at 65°C for 15 minutes as determined by agarose gel electrophoresis. | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of BstEII digested Lambda DNA and a minimum of 100 units of Taql Methyltransferase incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by  | Pass           |



M0219S / Lot: 10225044

Page 1 of 2



| Assay Name/Specification     | Lot # 10225044 |
|------------------------------|----------------|
| agarose gel electrophoresis. |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Nicele Castagnozzi Production Scientist

25 Jan 2024

Michael Tonello

Packaging Quality Control Inspector

26 Jan 2024