240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: GpC Methyltransferase (M.CviPI)

Catalog #: M0227S/L
Concentration: 4,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg Lambda DNA in 1 hour at 37°C in a total reaction

volume of 10 µl against cleavage by HaeIII restriction endonuclease.

 Lot #:
 0021709

 Assay Date:
 09/2017

 Expiration Date:
 09/2019

 Storage Temp:
 -20°C

Storage Conditions: 15 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-M0227S/L v1.0
Effective Date: 21 May 2018

Assay Name/Specification (minimum release criteria)	Lot #0021709
Endonuclease Activity (Nicking) - A 50 μl reaction in GC Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 40 units of GpC Methyltransferase (M.CviPI) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in GC Reaction Buffer containing 1 μg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 40 units of GpC Methyltransferase (M.CviPI) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (Methyltransferase) - A 20 μ l reaction in GC Reaction Buffer supplemented with 160 μ M SAM containing 1 μ g of Lambda DNA and 1 unit of GpC Methyltransferase (M.CviPI) incubated for 1 hour at 37°C followed by heat inactivation results in \geq 95% protection from digestion with 10 units of HaeIII in NEBuffer 2 incubated at 37°C for 1 hour as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in GC Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 40 units of GpC Methyltransferase (M.CviPI) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

Authorized by Derek Robinson 21 May 2018

nqa.
ISO 9001
Registered
Quality





Inspected by Timothy Meixsell 22 Sep 2017

Zimothy Meupelf