

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	T7 Exonuclease
Catalog Number:	M0263S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to produce 1 nmol of acid-soluble deoxyribonucleotide in a total reaction volume of 50 μl in 30 minutes at 37°C in 1X NEBuffer 4 with 0.15 mM sonicated duplex [ ³H]-DNA.
Packaging Lot Number:	10163517
Expiration Date:	08/2024
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl, 5 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 8.0 @ 25°C)
Specification Version:	PS-M0263S/L v1.0

T7 Exonuclease Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0263SVIAL	T7 Exonuclease	10160868	Pass	
B7004SVIAL	NEBuffer™ 4	10133928	Pass	

Assay Name/Specification	Lot # 10163517
<b>Endonuclease Activity (Nicking)</b> A 50 $\mu$ I reaction in NEBuffer 4 containing 1 $\mu$ g of supercoiled PhiX174 DNA and a minimum of 100 units of T7 Exonuclease incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b> T7 Exonuclease is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 10 units of T7 Exonuclease incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10163517
and a minimum of 10 units of T7 Exonuclease is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel	
electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

m J. Loei

John Greci Production Scientist 07 Sep 2022

Michae m. 11

Michael Tonello Packaging Quality Control Inspector 07 Sep 2022

