

New England Biolabs Certificate of Analysis

Product Name: Multiplex PCR 5X Master Mix
Catalog Number: M0284S
Concentration: 5 X Concentrate
Packaging Lot Number: 10122832
Expiration Date: 09/2023
Storage Temperature: -20°C
Specification Version: PS-M0284S v2.0
Composition (1X): 20 mM Tris-HCl (pH 8.9 @ 25°C), 50 mM KCl, 30 mM NH₄Cl, 2.5 mM MgCl₂, 0.3 mM dATP, 0.3 mM dCTP, 0.3 mM dGTP, 0.3 mM dTTP, 3.2 % Glycerol, 0.08 % IGEPAL® CA-630, 0.07 % Tween® 20, 67 units/ml Taq DNA Polymerase

Multiplex PCR 5X Master Mix Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0284SVIAL	Multiplex PCR 5X Master Mix	10122300	Pass

Assay Name/Specification	Lot # 10122832
PCR Amplification (15-plex PCR, Master Mix) A 25 µl reaction in 1X Multiplex PCR Master Mix and 0.15 µM primer mix containing 10 ng Human Genomic DNA for 35 cycles of PCR amplification results in the expected 15 products.	Pass
Protein Purity Assay (SDS-PAGE) Taq DNA Polymerase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Taq DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Multiplex PCR 5X Master Mix is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass


Assay Name/Specification	Lot # 10122832
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Taq DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 μl reaction in ThermoPol® Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 20 units of Taq DNA Polymerase incubated for 4 hours at 37°C and 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 μl reaction in ThermoPol® Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of Taq DNA Polymerase incubated for 30 minutes at 37°C and 75°C yields <10% degradation as determined by capillary electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 hour, Buffer) A 50 μl reaction in 2X Multiplex PCR Master Mix containing 1 μg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
17 Nov 2021



Michael Tonello
Packaging Quality Control Inspector
17 Nov 2021