

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	DNase I (RNase-free)
Catalog Number:	M0303L
Concentration:	2,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme which will completely degrade 1 µg of pBR322 DNA in 10 minutes at 37°C in DNase I Reaction Buffer. Complete degradation is defined as the reduction of the majority of DNA fragments to tetranucleotides or smaller.
Packaging Lot Number:	10234125
Expiration Date:	01/2026
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl (pH 7.6), 2 mM CaCl2 , 50 % Glycerol
Specification Version:	PS-M0303S/L v1.0

DNase I (RNase-free) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0303LVIAL	DNase I (RNase-free)	10213691	Pass	
B0303SVIAL	DNase I Reaction Buffer	10225938	Pass	

Assay Name/Specification	Lot # 10234125
Protein Purity Assay (SDS-PAGE) DNase I (RNase-free) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 2 units of DNase I (RNase-free) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 2 units of DNase I (RNase-free) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
<b>RNase Activity (ds RNA)</b> A 50 μl reaction in DNase I Reaction Buffer containing 10 μg of a dsRNA Ladder and a	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10234125
minimum of 100 units of DNase I (RNase-free) is incubated at 37°C. After incubation	
for 4 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Heid Mm

Heidi Church Production Scientist 14 Feb 2024

Michael Tr. W.

Michael Tonello Packaging Quality Control Inspector 03 Apr 2024

