

## New England Biolabs Certificate of Analysis

**Product Name:** *RNase Inhibitor, Human Placenta*  
**Catalog Number:** M0307L  
**Concentration:** 40,000 U/ml  
**Unit Definition:** One unit is defined as the amount of RNase Inhibitor, Human Placenta required to inhibit the activity of 5 ng of RNase A by 50%. Activity is measured by the inhibition of hydrolysis of cytidine 2', 3'-cyclic monophosphate by RNase A.  
**Packaging Lot Number:** 10106710  
**Expiration Date:** 06/2023  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM KCl, 20 mM HEPES (pH 7.6), 8 mM DTT, 50 % Glycerol  
**Specification Version:** PS-M0307S/L v1.0

| RNase Inhibitor, Human Placenta Component List |                                 |            |                      |
|--|---------------------------------|------------|----------------------|
| NEB Part Number                                | Component Description           | Lot Number | Individual QC Result |
| M0307LVIAL                                     | RNase Inhibitor, Human Placenta | 10109062   | Pass                 |

| Assay Name/Specification   | Lot # 10106710 |
|--|----------------|
| <b>RNase Activity (Extended Digestion)</b><br>A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of RNase Inhibitor, Human Placenta is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.                      | Pass           |
| <b>Protein Purity Assay (SDS-PAGE)</b><br>RNase Inhibitor, Human Placenta is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.   | Pass           |
| <b>Latent RNase Activity (Extended Digest)</b><br>A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 40 units of heat inactivated RNase Inhibitor, Human Placenta is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass           |
| <b>Exonuclease Activity (Radioactivity Release)</b><br>A 50 µl reaction in NEBuffer 4 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of RNase Inhibitor,  | Pass           |

| Assay Name/Specification  | Lot # 10106710     |
|---|--------------------|
| <p>Human Placenta incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p> <p><b>Endonuclease Activity (Nicking)</b><br/>A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 40 units of RNase Inhibitor, Human Placenta incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p> | <p><b>Pass</b></p> |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Bhairavi Jani  
Production Scientist  
22 Jun 2021



Michael Tonello  
Packaging Quality Control Inspector  
22 Jun 2021