

be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Human Alkyladenine Glycosylase (hAAG)
Catalog Number:	M0313S
Concentration:	10,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to create an AP site from 1 pmol of a 34-mer oligonucleotide duplex containing a single deoxyinosine site in a total reaction volume of 10 μ l in 1 hour at 37°C.
Packaging Lot Number:	10165607
Expiration Date:	10/2024
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.5 % Tween® 20 , 0.5 % IGEPAL® CA-630 , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-M0313S/L v1.0

Human Alkyladenine Glycosylase (hAAG) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0313SVIAL	Human Alkyladenine Glycosylase (hAAG)	10165606	Pass	
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10165338	Pass	

Assay Name/Specification	Lot # 10165607
Protein Purity Assay (SDS-PAGE) hAAG is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of hAAG incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of hAAG incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled	Pass





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10165607
PhiX174 DNA and a minimum of 100 units of hAAG incubated for 4 hours at 37°C results	
in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Lauren Higgins

Lauren Sears Higgins Production Scientist 13 Oct 2022

Michae 11.1

Michael Tonello Packaging Quality Control Inspector 02 Nov 2022

