240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Bst DNA Polymerase, Full Length

Catalog #: M0328S/L
Concentration: 5,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes

at 65°C.

 Lot #:
 0051803

 Assay Date:
 03/2018

 Expiration Date:
 3/2020

 Storage Temp:
 -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % Triton®X-100, 50 % Glycerol, (pH 7.1 @ 25°C)

Specification Version: PS-M0328S/L v1.0

Effective Date: 16 Oct 2015

Assay Name/Specification (minimum release criteria)	Lot #0051803
Endonuclease Activity (Nicking) - A 50 μ l reaction in ThermoPol® Reaction Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 50 units of <i>Bst</i> DNA Polymerase, Full Length incubated for 4 hours at either 37°C or 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of <i>Bst</i> DNA Polymerase, Full Length incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Phosphatase Activity (pNPP) - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 100 units <i>Bst</i> DNA Polymerase, Full Length incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) - <i>Bst</i> DNA Polymerase, Full Length is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 5 units of <i>Bst</i> DNA Polymerase, Full Length is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.	Pass







Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0051803
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of <i>Bst</i> DNA Polymerase, Full Length is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

Authorized by Melanie Fortier 16 Oct 2015







Inspected by
Tony Spear-Alfonso

27 Mar 2018