

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Bsu DNA Polymerase, Large Fragment
Catalog #:	M0330S/L
Concentration:	5,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 37°C.
<i>Lot</i> #:	0041703
Assay Date:	03/2017
Expiration Date:	03/2019
Storage Temp:	-20°C
Storage Conditions:	25 mM Tris-HCl , 50 mM NaCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-M0330S/L v1.0
Effective Date:	16 Oct 2015

Assay Name/Specification (minimum release criteria)	Lot #0041703
Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 50 units of <i>Bsu</i> DNA Polymerase, Large Fragment incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 50 units of <i>Bsu</i> DNA Polymerase, Large Fragment incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of <i>Bsu</i> DNA Polymerase, Large Fragment incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Phosphatase Activity (pNPP) - A 200 μ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl ₂ containing 2.5 mM <i>p</i> -Nitrophenyl Phosphate (pNPP) and a minimum of 100 units <i>Bsu</i> DNA Polymerase, Large Fragment incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) - <i>Bsu</i> DNA Polymerase, Large Fragment is \ge 97% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass



M0330S/L Lot: 0041703 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0041703
qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 5 units of <i>Bsu</i> DNA Polymerase, Large Fragment is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.	Pass
RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single- stranded RNA and a minimum of 1 μ l of <i>Bsu</i> DNA Polymerase, Large Fragment is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Single Stranded DNase Activity (FAM-Labeled Oligo) - A 50 µl reaction in NEBuffer 2 containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 50 units of <i>Bsu</i> DNA Polymerase, Large Fragment incubated for 30 minutes at 37°C yields <10% degradation as determined by capillary electrophoresis.	Pass

Authorized by Melanie Fortier 16 Oct 2015



UU. Int

Inspected by Tony Spear-Alfonso 11 Aug 2017