

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Exonuclease V (RecBCD)

Catalog Number: M0345L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to produce 1

nmol of acid-soluble deoxyribonucleotide from double-stranded DNA in

30 minutes at 37°C in a total reaction volume of 50 µl.

Packaging Lot Number: 10105432
Expiration Date: 01/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.1% Triton®X-100, (pH 7.5 @ 25°C)

Specification Version: PS-M0345S/L v1.0

Exonuclease V (RecBCD) Component List				
<b>NEB Part Number</b>	<b>Component Description</b>	Lot Number	Individual QC Result	
P0756SVIAL	Adenosine 5'-Triphosphate (ATP)	10094836	Pass	
M0345LVIAL	Exonuclease V (RecBCD)	10092695	Pass	
B7004SVIAL	NEBuffer™ 4	10101326	Pass	

Assay Name/Specification	Lot # 10105432
Endonuclease Activity (Nicked Double-Stranded DNA) A 50 μl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 μg of nicked PhiX174 RF II DNA and a minimum of 50 units of Exonuclease V (RecBCD) incubated for 4 hours at 37°C results in <10% loss in PhiX174 RF II DNA as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 100 units of Exonuclease V (RecBCD) incubated for 4 hours at 37°C results in <10% loss in supercoiled DNA as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Exonuclease V (RecBCD) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
RNase Activity (Extended Digestion)	Pass



M0345L / Lot: 10105432

Page 1 of 2

Assay Name/Specification	Lot # 10105432
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA	
and a minimum of 10 units of Exonuclease V (RecBCD) is incubated at 37°C. After	
incubation for 4 hours, >90% of the substrate RNA remains intact as determined by	
gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

John Greci Production Scientist

27 Apr 2021

Michael Tonello

Packaging Quality Control Inspector

27 Apr 2021

