

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Exonuclease V (RecBCD)

Catalog Number: M0345L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to produce 1

nmol of acid-soluble deoxyribonucleotide from double-stranded DNA in

30 minutes at 37°C in a total reaction volume of 50 µl.

Packaging Lot Number: 10137208
Expiration Date: 01/2024
Storage Temperature: -20°C

Storage Conditions: 50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.1% Triton®X-100, (pH 7.5 @ 25°C)

Specification Version: PS-M0345S/L v1.0

Exonuclease V (RecBCD) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
P0756SVIAL	Adenosine 5'-Triphosphate (ATP)	10121427	Pass	
M0345LVIAL	Exonuclease V (RecBCD)	10137205	Pass	
B7004SVIAL	NEBuffer™ 4	10133928	Pass	

Assay Name/Specification	Lot # 10137208
Protein Purity Assay (SDS-PAGE)	Pass
Exonuclease V (RecBCD) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of Exonuclease V (RecBCD) is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicked Double-Stranded DNA) A 50 µl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 µg of nicked PhiX174 RF II DNA and a minimum of 50 units of Exonuclease V (RecBCD) incubated for 4 hours at 37°C results in <10% loss in PhiX174 RF II DNA as determined by agarose gel electrophoresis.	Pass
Endonuclease Activity (Nicking)	Pass



M0345L / Lot: 10137208

Page 1 of 2

Assay Name/Specification	Lot # 10137208
A 50 µl reaction in NEBuffer 4 supplemented with 1 mM ATP containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 100 units of Exonuclease V (RecBCD)	
incubated for 4 hours at 37°C results in <10% loss in supercoiled DNA as determined	
by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

John Greci Production Scientist

Production Scientist 09 Feb 2022

Michael Tonello

Packaging Quality Control Inspector

09 Feb 2022