

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: RNA 5' Pyrophosphohydrolase (RppH)

Catalog Number: M0356S
Concentration: 5,000 U/ml

Unit Definition: One unit is the amount of enzyme that converts 1 µg 300 mer RNA

transcript into a XRN-1 digestible RNA in 30 minutes at 37°C.

Lot Number: 10013504
Expiration Date: 12/2020
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 20 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.01% Triton®X-100, (pH 7.5 @ 25°C)

Specification Version: PS-M0356S v1.0

RNA 5' Pyrophosphohydrolase (RppH) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0356SVIAL	RNA 5' Pyrophosphohydrolase (RppH)	10028256	Pass	
B7002SVIAL	NEBuffer™ 2	10023097	Pass	

Assay Name/Specification	Lot # 10013504
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 25 units of RNA 5' Pyrophosphohydrolase (RppH) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 25 units of RNA 5' Pyrophosphohydrolase (RppH) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 25 units of RNA 5' Pyrophosphohydrolase (RppH) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE)	Pass



M0356S / Lot: 10013504

Page 1 of 2

Assay Name/Specification	Lot # 10013504
RNA 5' Pyrophosphohydrolase (RppH) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 5 units of RNA 5' Pyrophosphohydrolase (RppH) is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Tony Spear-Alfonso Production Scientist

05 Oct 2018

Michael Tonello

Packaging Quality Control Inspector

13 Dec 2018

