

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: RNA 5' Pyrophosphohydrolase (RppH)

Catalog Number: M0356S
Concentration: 5,000 U/ml

Unit Definition: One unit is the amount of enzyme that converts 1 µg 300 mer RNA

transcript into a XRN-1 digestible RNA in 30 minutes at 37°C.

Packaging Lot Number: 10106683
Expiration Date: 08/2022
Storage Temperature: -20°C

Storage Conditions: 200 mM NaCl, 20 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

0.01% Triton®X-100, (pH 7.5 @ 25°C)

Specification Version: PS-M0356S v1.0

RNA 5' Pyrophosphohydrolase (RppH) Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0356SVIAL	RNA 5' Pyrophosphohydrolase (RppH)	10081895	Pass	
B7002SVIAL	NEBuffer™ 2	10102430	Pass	

Assay Name/Specification	Lot # 10106683
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 25 units of RNA 5' Pyrophosphohydrolase (RppH) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) RNA 5' Pyrophosphohydrolase (RppH) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 25 units of RNA 5' Pyrophosphohydrolase (RppH) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 25 units of RNA 5' Pyrophosphohydrolase (RppH) incubated for 4 hours at	Pass



M0356S / Lot: 10106683

Page 1 of 2

Assay Name/Specification	Lot # 10106683
37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	
RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 5 units of RNA 5' Pyrophosphohydrolase (RppH) is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani **Production Scientist**

04 May 2021

Josh Hersey

Packaging Quality Control Inspector 04 May 2021



M0356S / Lot: 10106683

Page 2 of 2