

## New England Biolabs Certificate of Analysis


**Product Name:** *beta-Agarase I*  
**Catalog Number:** M0392S  
**Concentration:** 1,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 200  $\mu$ l of molten low melting point or NuSieve agarose to nonprecipitable neoagaro-oligosaccharides in 1 hour at 42°C  
**Lot Number:** 10050911  
**Expiration Date:** 08/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 50 mM Bis-Tris-HCl, 1 mM EDTA, 50 % Glycerol, (pH 6.5 @ 25°C)  
**Specification Version:** PS-M0392S/L v1.0

| beta-Agarase I Component List |                                    |            |                      |
|-------------------------------|------------------------------------|------------|----------------------|
| NEB Part Number               | Component Description              | Lot Number | Individual QC Result |
| M0392SVIAL                    | $\beta$ -Agarase I                 | 10050910   | Pass                 |
| B0392SVIAL                    | $\beta$ -Agarase I Reaction Buffer | 10054316   | Pass                 |

| Assay Name/Specification                                                                                                                                                                                                                                                                                                                                                    | Lot # 10050911 |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <p><b>Endonuclease Activity (Nicking)</b><br/>           A 50 <math>\mu</math>l reaction in CutSmart® Buffer containing 1 <math>\mu</math>g of supercoiled PhiX174 DNA and a minimum of 1 unit of <math>\beta</math>-Agarase I incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>             | Pass           |
| <p><b>Exonuclease Activity (Radioactivity Release)</b><br/>           A 50 <math>\mu</math>l reaction in CutSmart® Buffer containing 1 <math>\mu</math>g of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 5 units of <math>\beta</math>-Agarase I incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>    | Pass           |
| <p><b>Non-Specific DNase Activity (16 Hour)</b><br/>           A 50 <math>\mu</math>l reaction in CutSmart® Buffer containing 1 <math>\mu</math>g of Lambda DNA and a minimum of 10 units of <math>\beta</math>-Agarase I incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | Pass           |
| <p><b>Protein Purity Assay (SDS-PAGE)</b><br/> <math>\beta</math>-Agarase I is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>                                                                                                                                                                                            | Pass           |

| Assay Name/Specification                                                                                                                                                                                                                                                                                                                        | Lot # 10050911     |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------|
| <p><b>RNase Activity (Extended Digestion)</b><br/>A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of β-Agarase I is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | <p><b>Pass</b></p> |

This product has been tested and shown to be in compliance with all specifications.



Ana Egana  
Production Scientist  
06 Sep 2019



Michael Tonello  
Packaging Quality Control Inspector  
06 Sep 2019