

New England Biolabs Certificate of Analysis

Product Name: LongAmp® Hot Start Taq 2X Master Mix
Catalog Number: M0533S
Concentration: 2 X Concentrate
Packaging Lot Number: 10236674
Expiration Date: 07/2025
Storage Temperature: -20°C
Specification Version: PS-M0533S/L v2.0
Composition (1X): 60 mM Tris-SO4 (pH 9.1 @ 25°C), 20 mM (NH4)2SO4, 2 mM MgSO4, 0.3 mM dATP, 0.3 mM dCTP, 0.3 mM dGTP, 0.3 mM dTTP, 3 % Glycerol, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 125 units/ml LongAmp® Hot Start Taq DNA Polymerase


LongAmp® Hot Start Taq 2X Master Mix Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0533SVIAL	LongAmp® Hot Start Taq 2X Master Mix	10226666	Pass

Assay Name/Specification	Lot # 10236674
Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation) A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200 µM dNTPs including [³ H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 10 units of LongAmp® Hot Start Taq DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 1X LongAmp® Hot Start Taq Master Mix containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (30 kb Human Genomic DNA, Master Mix) A 25 µl reaction in 1X LongAmp® Hot Start Taq Master Mix and 0.4 µM primers containing 500 ng Human Genomic DNA for 28 cycles of PCR amplification results in the expected 30 kb product.	Pass
PCR Amplification (30 kb Lambda DNA, Master Mix) A 25 µl reaction in 1X LongAmp® Hot Start Taq Master Mix and 0.4 µM primers containing 1 ng Lambda DNA for 28 cycles of PCR amplification results in the expected 30 kb product.	Pass

Assay Name/Specification	Lot # 10236674
<p>PCR Amplification (Hot Start, Human Genomic DNA, Master Mix) A 50 µl reaction in 1X LongAmp® Hot Start Taq Master Mix and 0.2 µM primers containing 2 ng Human Genomic DNA for 35 cycles of PCR amplification results in the expected 306 bp product and a decrease in non-specific genomic bands after pre-incubation at room temperature for 1 hour, when compared to a non-hot start control reaction.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of LongAmp® Hot Start Taq 2X Master Mix is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 2.5 units of LongAmp® Hot Start Taq DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.


Lea Antonopoulos
Production Scientist
07 Mar 2024


Michael Tonello
Packaging Quality Control Inspector
16 Apr 2024