

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

pBR322 DNA-MspI Digest
N3032S/L
1 mg/ml
N/A
1561804
04/2018
4/2020
-20°C
10 mM Tris-HCl (pH 8.0), 1 mM EDTA
PS-N3032S/L v1.0
30 Mar 2017

Assay Name/Specification (minimum release criteria)	Lot #1561804
A260/A280 Assay - The ratio of UV absorption of pBR322 DNA-MspI Digest at 260 and 280 nm is between 1.8 and 2.0.	Pass
DNA Concentration (A260) - The concentration of pBR322 DNA-MspI Digest is between 1000 and 1050 μ g/ml as determined by UV absorption at 260 nm.	Pass
Electrophoretic Pattern (Marker) - The banding pattern of pBR322 DNA-MspI Digest on a 3% agarose gel shows discrete, clearly identifiable bands at each band of the marker, when stained with Ethidium Bromide at a concentration of 0.5 μ g/ml.	Pass
Non-Specific DNase Activity (DNA, 16 hour) - A 50 µl reaction in 1X NEBuffer 2 containing 5 µg of pBR322 DNA-MspI Digest incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

otunen

Authorized by Derek Robinson 30 Mar 2017



Mulhiceheter

Inspected by Vanessa Mathieu-Sheltry 17 Apr 2018